

Emergency Department

Emergency Care Screening

Common Screenings done in the ED

- Stroke
 - FAST Exam (facial drooping, arm weakness, speech difficulties, time)
- Depression
 - PHQ-9
- Domestic Violence
 - SAFE Questions (Stress/Safety, Afraid/Abused, Friends/Family, Emergency plan)
- Alcohol/Drug Abuse
 - SBIRT
 - (NIAAA: 1) How many days per week do you drink alcohol?, On a typical day when you drink, how many drinks do you have?, 3) what is the maximum number of drinks you had on any given day in the past month?))
 - [NIDA_Modified_ASSIST_prescreen.pdf](#)
 - DAST [DAST.pdf](#)
- Tobacco Use
 - Y/N, form, amount, length of use, age started
- Communicable diseases
 - Influenza - flu shot received, Y/N
 - Ebola - travel outside of US
 - Zika - ??
- Seatbelt Use
 - Y/N, always, sometimes, never...
- Fall Risk
 - Fall within 12 months, number, last
- SDoH
 - PREPARE
 - SEEK

(note to Laura - build the table to see how this data fits into the resources (questionnaire, questionnaire response, who is asking the question, when, what is done with the results/content/answers received)

Preamble: The use case focus for this track will be limited to the screening done during the patient's ED visit.

Emergency Care Triage

1. Patient Name
 - a. Mrs. Patricia (Pat) Chess
 - b. 42-year-old full time office assistant, was involved in a road accident
2. Patient Condition
 - a. Post MVA
 - i. **Pat is admitted to ER**, treated and transferred to OR (see primary story)
 - ii. Diagnostic abdominal, peripheral, and chest and spinal X-rays and CT, showed a closed undisplaced comminuted fracture left mid shaft femur, fracture left ribs 6-7 with spleen and liver contusions. No fracture vertebrae detected.

Emergency Care Admission

Emergency Care Care Provision and Treatment

Emergency Care Discharge